

**COMPOSITION:** Each 500 mL of sterile aqueous solution contains:

|   |        |
|---|--------|
| Calcium .....   | 10.8 g |
| (as calcium borogluconate, equivalent to calcium gluconate 23.2%) |        |
| Potassium .....   | 8.0 g  |
| (as potassium chloride)   |        |
| Phosphorus .....  | 2.5 g  |
| (as sodium hypophosphite • H <sub>2</sub> O)                      |        |
| Magnesium .....   | 1.6 g  |
| (as magnesium borogluconate)                                      |        |
| Dextrose • H <sub>2</sub> O .....                                 | 75.0 g |

**Milliequivalents per liter**

**Cations**

|                 |             |
|-----------------|-------------|
| Calcium .....   | 1,080 mEq/L |
| Potassium ..... | 410 mEq/L   |
| Magnesium ..... | 261 mEq/L   |
| Sodium .....    | 161 mEq/L   |

**Anions**

|                     |             |
|---------------------|-------------|
| Borogluconate ..... | 1,341 mEq/L |
| Chloride .....      | 410 mEq/L   |
| Hypophosphite ..... | 161 mEq/L   |

Store between 15°C and 30°C (59°F and 86°F).

18-805

RMS 92-543

Rev. 03-14

Lot No.

Exp. Date

NDC 57319-544-07

# Cal-Phos #2 With Potassium Injection

For Treatment of Milk Fever in Cattle

**FOR ANIMAL USE ONLY**

**KEEP OUT OF REACH OF CHILDREN**

**Net Contents: 500 mL**

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**PHOENIX™**

Manufactured for:  
Clipper Distributing Company, LLC  
St. Joseph, MO 64507

**INDICATIONS:** For use as an aid in the treatment of hypocalcemia (parturient paresis, milk fever), hypomagnesemia (grass tetany), and other conditions associated with calcium, magnesium, phosphorus and potassium deficiencies in cattle.

**CONTRAINDICATIONS:** Do not administer this product to animals showing signs of cardiac distress.

**PRECAUTIONS:** Administration should be made slowly and with care to avoid adverse effects such as heart block or shock. Perivascular or subcutaneous deposition of hypertonic solutions may result in severe inflammation at the injection site.

**CAUTION:** This product contains no preservatives. Use entire contents when first opened. Discard any unused solution.

**DOSAGE AND ADMINISTRATION:**

Warm solution to body temperature. The usual intravenous dose in cattle is 500 mL per 800 to 1,000 pounds of body weight.

TAKE TIME



OBSERVE LABEL DIRECTIONS

Trademarks are property of  
Clipper Distributing Company, LLC

Manufactured by  
Nova-Tech Inc.  
Grand Island, NE 68801



3