

**Dosage and Administration:** Dosages should be restricted to the lowest level necessary to control clinical signs. If no therapeutic response is noted in seven days, the diagnosis should be redetermined.

**Recommended Dose:** 2.5 mg per 10 lb (4.5 kg) body weight per day in divided doses six to ten hours apart.

**WARNING:** Clinical and experimental data have demonstrated that corticosteroids administered orally or parenterally to animals may induce the first stage of parturition when administered during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta and metritis. Additionally, corticosteroids administered to dogs, rabbits and rodents during pregnancy have resulted in cleft palate in offspring. Corticosteroids administered to dogs during pregnancy have also resulted in other congenital anomalies, including deformed forelegs, phocomelia and anasarca.

Lot No.

Exp. Date

NDC 57319-441-16

# PrednisTab<sup>®</sup>

(prednisolone tablets), USP

## 5 mg

**For Oral Use in Dogs Only**

Approved by FDA under NADA # 140-921

**FOR ANIMAL USE ONLY**  
**KEEP OUT OF REACH**  
**OF CHILDREN**

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**PHOENIX**  
Manufactured for:  
Clipper Distributing Company, LLC  
St. Joseph, MO 64507

**Net Contents:**  
**1000 Tablets**

**Each Tablet Contains:**

Prednisolone, USP ..... 5 mg

**Important:** Read the accompanying leaflet carefully and completely before use.

**KEEP OUT OF REACH OF CHILDREN**

**STORAGE:** Store at controlled room temperature 20° - 25°C (68° - 77°F), excursions permitted between 15° - 30°C (59° - 86°F).

TAKE TIME



OBSERVE LABEL DIRECTIONS

Trademarks are property of  
Clipper Distributing Company, LLC

Manufactured by:  
LLOYD, Inc.  
Shenandoah, IA 51601

Rev. 09-19

