

INDICATIONS: For the correction of electrolyte depletion, metabolic acidosis and dehydration of cattle, calves, horses, sheep and swine.

DOSAGE AND ADMINISTRATION: May be injected intravenously, subcutaneously or intraperitoneally (except in horses) using strict aseptic technique.

Cattle and Horses: 2 to 5 mL per pound of body weight depending on size and condition of animal, repeated 1 to 3 times daily or as needed.

Swine and Sheep: 2 to 5 mL per pound of body weight depending on size and condition of animal, repeated 1 to 3 times daily or as needed.

If administered subcutaneously divide the dosage into several sites of injection and massage points of injection to aid in absorption and help prevent inflammation and/or sloughing.

Store between 15°C and 30°C (59°F and 86°F)

TAKE TIME



OBSERVE LABEL
DIRECTIONS

RMS 92-541
18-803-60

Rev. 06-10

NDC 57319-545-08

Lactated Ringers Injection

Sterile Nonpyrogenic Solution

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

Net Contents: 1000 mL

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.



Manufactured for:
Clipper Distributing Company, LLC
St. Joseph, MO 64507

Each 100 mL contains:
Sodium Chloride 600 mg
Sodium Lactate 310 mg
Potassium Chloride 30 mg
Calcium Chloride Dihydrate 20 mg
Water for Injection q.s.

The Calcium, Potassium and Sodium contents are approximately 2.7, 4.0, and 130 mEq/liter, respectively. Total Osmolar Concentration: 269 mOsm per liter (calculated).

CAUTION: Solution should be warmed to body temperature prior to administration and administered at a slow rate. This is a single dose unit. It contains no preservatives. Use entire contents when first opened.

WARNING: Do not administer to horses by intraperitoneal injection. Do not administer to animals with inadequate renal function. Not for use in lactic acidosis.

Manufactured by
Nova-Tech Inc.
Grand Island, NE 68801

Trademarks are property of
Clipper Distributing Company, LLC

Lot No.

Exp. Date

