

INDICATIONS: Lidocaine is a potent local anesthetic for producing epidural and nerve conduction anesthesia.

CONTRAINDICATIONS: Lidocaine is contraindicated in animals with a known hypersensitivity to the drug.

PRECAUTIONS: Lidocaine is usually well tolerated. Nevertheless, as with all local anesthetics, untoward effects may occur due to hypersensitivity, faulty technique, overdosage and inadvertent intravascular or subarachnoid injection. In case of respiratory arrest, immediate resuscitation with oxygen is indicated.

DOSAGE AND ADMINISTRATION:

Epidural

Cattle and Horses - 5 to 15 mL.

Dogs and Cats - 1 mL per 10 pounds of body weight.

Nerve Block

Cattle and Horses - 5 to 20 mL.

L-2000-04

Lot No.

Exp. Date

Rev. 06-10

NDC 57319-533-05

Lidocaine Hydrochloride Injectable-2%

**FOR ANIMAL USE ONLY
KEEP OUT OF REACH OF CHILDREN**

Net Contents: 100 mL


PHOENIX™

Manufactured for:
Clipper Distributing Company, LLC
St. Joseph, MO 64507

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

COMPOSITION: Each mL of sterile aqueous solution contains:

Lidocaine Hydrochloride	2.0%
Propylene Glycol	5.2%
Sodium Chloride	0.5%
Sodium Lactate	0.5%

with Methylparaben 0.15%, Sodium Metabisulfate 0.10%, Propylparaben 0.03% and Disodium Edetate 0.001% as preservatives.

Store at controlled room temperature between 15°C and 30°C (59°F and 86°F).

TAKE TIME



OBSERVE LABEL
DIRECTIONS

Trademarks are property of
Clipper Distributing Company, LLC

Manufactured by
Sparhawk Laboratories, Inc.
Lenexa, KS 66215

