INDICATIONS: Lidocaine is a potent local anesthetic for producing epidural and nerve conduction anesthesia

CONTRAINDICATIONS: Lidocaine is contraindicated in animals with a known hypersensitivity to the drug.

PRECAUTIONS: Lidocaine is usually well tolerated. Nevertheless, as with all local anesthetics, untoward effects may occur due to hypersensitivity, faulty technique, overdosage and inadvertent intravascular or subarachnoid injection. In case of respiratory arrest, immediate resuscitation with oxygen is indicated

DOSAGE AND ADMINISTRATION:

Epidural

Cattle and Horses - 5 to 15 mL.

Dogs and **Cats -** 1 mL per 10 pounds of body weight.

Nerve Block

Cattle and Horses - 5 to 20 ml.

LOT NO.:

EXP. DATE:

NDC 57319-533-06

Lidocaine Hydrochloride Injectable-2%

FOR ANIMAL USE ONLY
KEEP OUT OF REACH OF CHILDREN

Net Contents: 250 mL

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Clip

PHOENIX™

Manufactured for: Clipper Distributing Company, LLC St. Joseph, MO 64507 **COMPOSITION:** Each mL of sterile aqueous solution contains:

with Methylparaben 0.15%, Sodium Metabisulfate 0.10%, Propylparaben 0.03% and Disodium Edetate 0.001% as preservatives.

Store at controlled room temperature between 15°C and 30°C (59°F and 86°F).

L-2000-05 Rev. 04-11

KE TIME OBSERVE DIRECTION

Trademarks are property of Clipper Distributing Company, LLC

Manufactured by Sparhawk Laboratories, Inc. Lenexa, KS 66215

