

INDICATIONS: For use as a supplemental nutritive source of vitamins A and D in cattle, sheep and swine.

CAUTION: Do not inject into meat animals within 60 days of marketing. Administration of this product to well nourished animals may cause hypervitaminosis D which may result in hypercalcemia. This condition may be harmful to the animal.

DOSAGE AND ADMINISTRATION: Inject intramuscularly or subcutaneously preferably in the neck area using aseptic technique.

Calves - 1/2 to 1 mL

Adult Cattle - 2 to 4 mL

Lambs - 1/4 to 1/2 mL

Adult Sheep - 1 to 2 mL

Weaning Pigs - 1/4 to 1/2 mL

Adult Swine - 1 to 2 mL

These suggested dosages may be repeated after 60 days, if necessary.

V-0332-05

Rev. 4-04

TAKE TIME



OBSERVE LABEL DIRECTIONS

Lot No.

Exp. Date

NDC 57319-424-06

Vitamin A D Injection

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**Net Contents:
250 mL**

Manufactured For:

PHOENIX
PHARMACEUTICAL, INC.
St. Joseph, MO 64503

COMPOSITION: Each mL of sterile solution contains:

Vitamin A.....500,000 IU

Vitamin D₃.....75,000 IU

In an emulsifiable base with vitamin E

(antioxidant), N-methylpyrrolidone,

polyoxyethylated (30) castor oil,

propylene glycol dicaprylate,

polyoxyethylene (20) sorbitan

monooleate and benzyl alcohol 2%.

Store at controlled room temperature

between 15° and 30°C (59-86°F).

Protect from light and

excessive heat.



Manufactured by
Sparhawk Laboratories
Lenexa, KS 66215